

Insecticidal effects of essential oils from six aromatic and medicinal plants on the pine processionary caterpillar (*Thaumetopoea pityocampa* Schiff)

Foued Hasnaoui^{1*}, Ikbel Zouaoui^{1,3}, Sana Dallali^{1,2} and Faten Dibouba¹

¹Silvo-Pastoral Institute of Tabarka, University of Jendouba, Tabarka, Tunisia.

²Research Laboratory for Agricultural Production Systems and Sustainable Development, Higher School of Agriculture, 1121 Mograne, Zaghouan, Tunisia.

³Biology Department, Faculty of Sciences of Tunis, University of Tunis El Manar, 2092 Tunis, Tunisia.

Accepted 20 August, 2020

ABSTRACT

The objective of the present study is to test some essential oils in order to recommend a method of effective control of *Thaumetopoea pityocampa* Schiff. Essential oils composition, polyphenols content, and antioxidant activity of six aromatic and medicinal plants leaf from Tabarka were investigated. The essential oils obtained by steam distillation from six plant species naturally grown in Tabarka forests were tested at three different concentrations in order to evaluate their effectiveness against the larvae of pine processionary (*Thaumetopoea pityocampa* Schiff). Our results revealed the effect of *Pinus pinaster* essential oils which gave the highest mortality rate of 80.55% and the fastest on *Thaumetopoea pityocampa* Schiff. followed by *Lavandula stoechas* and *Eucalyptus camaldulensis* with 74.81 and 72.81%. The other species tested were *P. pinea*, *Juniperus oxycedrus* and *Cupressus sempervirens* also had a considerable effect on caterpillar mortality with percentages 66.66, 58.26 and 40.49%. Then, a phytochemical study of the six species tested was carried out which showed that some oils can be used as a spray substance to control *T. pityocampa* Schiff due to their compositions in active substances.

Keywords: Essential oils, pine processionary caterpillar (*Thaumetopoea pityocampa* Schiff), insecticidal activity, Tabarka forests.

*Corresponding author. E-mail: hasnaoui.foued@gmail.com.

INTRODUCTION

Tunisian pine forests are affected by various depredation phenomena (drought, anthropogenic activities, and insect pests). In addition to bark beetles, defoliation by the pine processionary (*Thaumetopoea pityocampa* Schiff) is highly important. Defoliation causes tree growth losses and makes them more susceptible to attack by secondary insects and diseases, (FAO, 2010). The continuous use of synthetic chemicals for insect control apart from their high efficacy and selectivity has raised concerns about their potential safety for human health and the environment. Secondary metabolites constitute a biologically and chemically interesting group of substances extracted from plants. Essential oils of plants show many biological activities in addition to their use in food, flavor, perfumery,

cosmetic and pharmaceutical industries as natural antioxidants (Wei and Shibamoto, 2010; Mothana et al., 2012). Essential oil has been used since ancient times for medicinal purposes and known for its anti-rheumatic, anti-inflammatory, and antispasmodic properties (Benincá et al., 2011; Zaouali et al., 2013). It has demonstrated powerful antimutagenic, antibacterial and chemopreventive properties (Okoh et al., 2010; Ayadi et al., 2011). The secondary metabolites grouped as essential oil impart them the much needed curative properties (Derwich et al., 2010).

Secondary metabolites from plants have important biological and pharmacological activities, such as anti-oxidative, anti-allergic, antibiotic, hypoglycemic, and anti-

carcinogenic (Stanković, 2011). The essential oils have different kinds of effects for insects: larvicides, autocides, repellents or growth inhibitors. Most essential oils act through disrupting the structure of the cell membranes. However, for some, neurotoxic effects have been demonstrated, due to interactions with neurotransmitters such as gamma-aminobutyric acid and octopamine or by inhibition of acetylcholinesterase. Some essential oils can potentiate the action of other molecules by inhibiting cytochromes P450 which normally taxed them. Because of their volatility and small size, many of the constituents of essential oils interact with the odor receptors of insects, triggering various behaviors: flight, attraction, oviposition, etc. (Tabassum and Vidyasagar, 2013).

The aim of this study was to find out whether essential oils from various plants might be used as insecticides for the control of *Thaumetopoea pityocampa* Schiff.

MATERIALS AND METHODS

Animal material

Estimating the health condition of studied pine forests and their infestation by insects in Tabarka shows that the pine processionary caterpillar is the greatest in the pine forests of Northwestern Tunisia.

To study the seasonal development of the pine processionary caterpillar and to test the effectiveness of its control, weekly visits to the study areas were carried out. (Figure 1)

During our study, the periods of different stages of the pine processionary caterpillar was refined (Figure 2).

It was recognized that the processional phase starts at the beginning of February and may last until the end of March.

Plant material

Leaves of some aromatic and medicinal plants (*Pinus pinaster*, *Lavandula stoechas*, *Eucalyptus camaldulensis*, *Pinus pinea*, *Juniperus oxycedrus*, *Cupressus sempervirens*) were collected from Tabarka forests, a preserved environment where the plants are vigorously producing very large biomass. The harvested materials were air-dried at room temperature ($20 \pm 2^\circ\text{C}$) for one week, ground, sieved through a 0.5 mm mesh screen to obtain a uniform particle size.

Total lipid extraction

Samples of ground powder (1g) in triplicate were weighed and extracted using the modified method of Bligh and Dyer (1959). A chloroform/methanol mixture (2:1, v/v) was used for total lipid extraction. The mixture was shaken and centrifuged (Eppendorf 5810R) at $3,000 \times g$ for 10 min to allow phase development. The organic layer containing total lipids was collected and filtered. The total extracted lipid material was recovered after the solvent was removed in a stream of nitrogen. The total fat content (TFC) was expressed as a percent of the dry matter and was calculated using the following formula (Gandour et al., 2011):

$$\text{TFC}(\% \text{ of dw}) = 100 \times \frac{m}{M} \quad (1)$$

Where TFC is the total fat content, m: the mass of extracted oil

(mg), M: the mass of dry matter (mg) and dw: the dry weight.

Essential oil isolation

The air-dried materials were hydro distilled for 3 h. The obtained distillate was extracted using diethyl-ether as solvent (v/v) and dried over anhydrous sulphate sodium. The organic layer was then concentrated at 35°C using a Vigreux column and the essential oil stored at -4°C prior to analysis. The essential oil yield was estimated according to dry leaves matter by using the following equation (Sangun et al., 2007):

$$\text{Rhe}(\%) = 100 \times \frac{M_{he}}{M_s} \quad (2)$$

Where M_{he} : essential oil mass (g), M_s : dry leaves matter mass (g), R_{he} : essential oil yield (%).

Essential oil gas chromatography analysis

Gas chromatography analyses were done with Shimadzu HRGC-2010 gas chromatograph equipped with flame ionisation detector (FID), Auto-injector and auto-sampler. An apolar column Rtx-1 (30 m \times 0.25 mm, 0.32 μm film thickness) was used. The oven temperature was held at 50°C for 10 min then programmed at $2^\circ\text{C}/\text{min}$ to 190°C then held isothermal for 10 min. The injector and detector temperature were programmed at 230°C . The flow of the carrier gas (N_2) was 1.6 ml/min and the split ration was 1:20. Injection volume for all samples was 0.5 μl of diluted oils in *n*-pentane. The volatile compounds were identified by comparison of their retention indices (RI) relative to (C_7 – C_{20}) *n*-alkanes with those of literature and/or with those of authentic compounds available in our laboratory. Relative percentage amounts of the identified compounds were obtained from the electronic integration of the FID peak areas.

Preparation of methanolic extracts

Methanolic extracts of some aromatic and medicinal plants were obtained, as described by Eghdami and Sadeghi (2010), with slight modification by Benhammou et al. (2009).

Total phenolic content

Total phenolics were determined with Folin–Ciocalteu (F-C) assay as per Singleton et al. (1999) method and slightly modified by Dewantoet al. (2002). The total phenolic content was expressed as mg of gallic acid equivalents per gram of dry weight (mg GAE/g DW) through the calibration curve of gallic acid.

Total flavonoids content

Total flavonoids content was determined according to the aluminium chloride colorimetric method (Djeridane et al., 2006). The total flavonoids content was calculated on the basis of the calibration curve of quercetin and expressed as mg quercetin equivalents per g dry weight (mgQE/g DW).

Antiradical scavenging activity by DPPH method

Plant extracts were tested for the scavenging effect on DPPH. This method is based on the measurement of the capacity of



Figure 1. The nest (left) and foliage damage by pine processionary caterpillars (right).




Figure 2. Cycle of *Thaumetopoea pityocampa*.

antioxidants to trap the 2,2-diphenyl-1-picrylhydrazil (DPPH) radical according to the method described by Braca et al. (2001). The DPPH radical scavenging activity in terms of percentage was calculated according to the following equation (Kumaran and Karunakaran, 2006):

$$I(\%) = \left[\frac{A_0 - A_1}{A_0} \right] \times 100 \quad (3)$$

Where I was DPPH inhibition (%), A_0 was the absorbance of the control, and A_1 was the absorbance of the extract/standard.

The concentration of the sample required for 50% inhibition was determined and represented as IC_{50} for each test solution which is expressed as $\mu\text{g/ml}$. All measurements were performed in triplicate.

Biological assays

Thaumetopoea pityocampa Schiff larvae were captured at the fourth and fifth instar, available on the Tabarka pines. The larvae were kept for use in studying the effect of certain oils on their behavior in laboratory. The essential oils were poured on 50 larvae, placed in Petri dishes (\varnothing 18 mm). The effects of all essential oils were evaluated by measuring the mortality time for killing 50 larvae (100% mortality). The complete test set was replicated four times for each essential oil. The average of four measurements was reported as mean mortality time (TEM).

Statistical analysis

All the analyses were performed in triplicate for each sample and expressed as mean \pm standard deviation (SD). One way analysis of variance (ANOVA) followed by Duncan's Multiple Range Test was applied to compare means at the significance level $p < 0.05$. All analyses were performed using SAS v 9.1 software package.

RESULTS AND DISCUSSION

Total lipid content

The amounts of total lipids in six aromatic and medicinal plants leaves are presented in Table 1.

Variation in the lipid content between species ($P > 0.0003$), of which *Eucalyptus* has the highest value of 6.64%.

In fact, Lavender, *Juniperus*, and *Cupressus* have close values (4.86, 5.29 and 4.92% respectively). However, *Pinus pinea* has the lowest value of 2.79%.

These variations in total lipids content among ecotypes of *Sesamum indicum* L. could be mainly due to the effect of genetic factors as well as environmental, edaphic, and cultural conditions (Uzun et al., 2002; Ravi et al., 2007).


Yields and essential oils components

Essential oil yields vary considerably among the six plants studied, *Lavandula stoechas*, *Cupressus sempervirens*, *Pinus pinaster*, *Pinus pinea*, *Juniperus oxycedrus*, *Eucalyptus camaldulensis*.

Indeed, the highest yield is recorded in *Cupressus* (0.25%) while *L. stoechas* (0.05%) has the lowest yield as shown in Figure 3. Variation in oil yield can be attributed to some factors such as conditions of plant growth, environment, and region. The oil yield during plant growth is particularly susceptible to environmental

Table 1. Total lipid content (%) of leaves of six aromatic and medicinal plants (%).

Species	Lipid content (%)
<i>Pinus pinea</i>	2.79 ± 0.23 ^d
<i>Lavandula stoechas</i>	4.86 ± 0.40 ^{bc}
<i>Pinus pinaster</i>	3.74 ± 0.10 ^{cd}
<i>Juniperus oxycedrus</i>	5.29 ± 1.45 ^b
<i>Cupressus sempervirens</i>	4.92 ± 0.58 ^{bc}
<i>Eucalyptus camaldulensis</i>	6.64 ± 0.26 ^a

**Figure 3.** Yield of essential oil (%).

conditions namely light, nutrient availability, day length, and daily temperature (Skoula et al., 2000; Msaada et al., 2009).

The essential oil compositions of Tunisian six aromatic and medicinal plants were determined by GC–FID analysis.

Compounds were identified with a significant variation of the essential oil chemical composition according to studied sites (Table 2).

The chemical class characterization of leaf essential oil showed the prevalence of monoterpene hydrocarbons. The oxygenated sesquiterpenes formed the second main class and the oxygenated monoterpenes formed the third main class.

The remaining fractions, such as aldehydes and sesquiterpene hydrocarbons, formed the minor essential oil chemical classes of six plant leaves studied.

This variability can be explained by the influence of various factors. For instance, it was previously shown for the majority of vegetal species, that primary and secondary metabolism are strongly influenced by plant physiology and genetic, climatic and edaphic factors, the harvest period, inducing changes in chemical qualitative and quantitative composition (Msaada et al., 2009; Hosni et al., 2011; Jemâa, 2014). Thus, the same species may show different chemical compositions from a region to

another due to genetic variability influenced by environmental factors (Pritchard et al., 2000). The influence of environmental factors in the chemical composition of essential oils has also been reported in *A. absinthium* (Bailen et al., 2013). Variations in the relative amounts of these compounds under different environmental conditions indicate their adaptively ecological value.

Thereby, the variations observed can be explained by the influence of the altitude. Essential oil composition of peppermint (*Mentha piperita*) was different because of the altitude and irregular daylight (Yazdani et al., 2002). Plants are grown and produced in different ecosystems and sites under the influence of different potential factors, including the altitude as one of the vital determinants in the quantity and quality of the plants (Mahdavi et al., 2013). Chemical components of the essential oils are cited to vary qualitatively and quantitatively according to geographical location and environmental conditions (Bakkali et al., 2008; Oliveira et al., 2013).

Kouyokhi et al. (2008) reported that phytochemical variations were found not only among samples from different regions but also among samples from the same region with different altitudes, reflecting the effect of environment on essential oil components. Habibi et al.

Table 2. Chemical composition of essential oils obtained from leaves of six aromatic and medicinal plants.

Volatile compound	Species					
	A	B	C	D	E	F
Hexanol	-	0.4	0.3	0.2	0.9	0.4
Nonane	0.1	-	0.4	0.3	0.6	0.3
α -Thujene	0.15	0.11	0.13	-	0.2	0.25
trans-2-Hexanal	0.18	0.22	-	0.19	0.3	0.4
(z)-2-heptenal	0.55	0.34	0.57	0.63	0.2	0.15
α -Pinene	7.2	6.8	3.35	4.3	2.6	1.9
Camphene	2.32	2.10	1.01	3.1	2.2	1.81
1-Octen-3-ol	-	3.2	4.1	3.1	0.6	0.9
Sabinene	-	-	0.1	1.9	1.1	2.1
β -Pinene	7.01	6.8	3.35	-	0.1	4.3
β -Myrcene	0.11	1.3	-	-	0.3	0.9
Decane	0.18	-	-	0.2	1.3	0.7
α -Terpinene	0.6	0.55	0.3	1.4	0.3	1.3
para-Cymene	-	0.2	1.2	0.7	-	-
γ -terpinene	1.9	1.39	1.75	0.63	3.4	-
para-Cymenene	-	-	-	0.3	0.2	1.3
Nonanal	0.4	0.9	1.3	5.6	4.9	0.2
α -Terpinolene	7.02	14.1	0.1	-	11	1.5
Undecane	1.6	-	-	-	0.3	-
cis-para-Menth-2-ene-1-ol	2.22	1.6	1.92	0.36	1.1	0.55
2-phenylethanol	-	-	0.3	1.8	1.4	-
trans-para-Menth-2-ene-1-ol	-	2.6	1.7	4.6	1.5	-
α -Terpineol	0.5	15.2	14.21	5.6	12	5.9
cis-Piperitol	-	0.9	1.6	2.9	6.7	-
Carvacrylmethyloxyde	0.14	0.11	0.58	0.09	0.2	0.78
Carvacrolmethylether	0.12	-	-	-	-	-
Linolylacetate	-	0.6	-	1.4	5	-
Linalylacetate	-	1.3	0.17	4.3	2.3	-
Perillaldehyde	1.5	2.3	2.9	-	-	1.6
Decanol	-	-	0.3	-	1.9	1.8
Thymol	-	4.5	7.1	-	0.3	14
Bornylacetate	-	1.6	8.9	-	-	7.9
Carvacrol	-	4.6	-	-	-	-
Pinocarvylacetate	2.6	-	1.9	-	1.6	-
Tridecane	1.7	-	-	-	0.9	-
α -Terpinylacetate	1.3	-	-	1.6	4.9	1.5
Nerylacetate	2.9	-	1.67	1.3	1.9	1.7
α -cubebene	1.58	-	-	11	1.9	18
Aromadendrene	1.9	-	1.7	0.55	7.3	2.3
β -Ionone	2.3	-	-	0.69	0.3	1.6
Germacrene D	-	1.4	-	-	0.9	1.9
γ -Cadinene	5.3	3	9	1.8	12	-
δ -Cadinene	-	-	-	0.6	1.3	-
α -Calacorene	-	-	-	-	-	-

(A) *Pinus pinea*, (B) *Lavandula stoechas*, (C) *Pinus pinaster*, (D) *Juniperus oxycedrus*, (E) *Cupressus sempervirens*, (F) *Eucalyptus camaldulensis*.

(2006) also reported that the essential oils of *Thymus kotschyanus* differ in different altitudes, but as the altitude increases, the amount of essential oil dropped.

Total phenolic content

Total phenolic values of six aromatic and medicinal plant

Table 3. Total phenolic and flavonoid content and DPPH radical-scavenging activity (IC₅₀) from the leaves of six aromatic and medicinal plants.

Species	Total phenolic content (mg GAE/g DW)	Total flavonoids content (mg QE/g DW)	IC ₅₀ (µg/ml)
<i>Pinus pinea</i>	119.73 ± 1.15 ^b	12.82 ± 19.24 ^a	1380.13 ± 1.00 ^b
<i>Lavandula stoechas</i>	86.12 ± 5.78 ^c	10.89 ± 1.33 ^b	1865.51 ± 0.57 ^a
<i>Pinus pinaster</i>	89.06 ± 2.00 ^c	5.48 ± 0.43 ^c	154.14 ± 1.00 ^c
<i>Juniperus oxycedrus</i>	156.53 ± 2.80 ^a	6.68 ± 0.29 ^d	27.72 ± 1.00 ^e
<i>Cupressus sempervirens</i>	64.61 ± 0.40 ^d	3.29 ± 0.30 ^e	30.42 ± 1.00 ^e
<i>Eucalyptus camaldulensis</i>	87.64 ± 2.01 ^c	6.14 ± 0.73 ^d	50.0 ± 10.0 ^d

Values of the same column with different superscripts (a–c) are significantly different at $p < 0.05$.

leaves are given in Table 3.

This variation can be attributed to several factors climatic and environment, geographical area, drought, diseases (Ebrahimi et al., 2008; Andarwulan et al., 2010), harvest time, and stage of plant development (Miliauskas et al., 2004).

In addition, genetic factors and growing conditions may play a crucial role in the composition of secondary metabolites, including phenolic acids (Islam et al., 2003; Hashempour, et al., 2010). The method of extraction and quantification also influences the estimation of total phenols content (Lee et al., 2003). Phenols are very important plant constituents not only because of their scavenging ability on free radicals due to their hydroxyl groups. Therefore, the phenolic content of plants may contribute directly to their antioxidant action and it is likely that the activity of the extracts is due to these compounds (Tosun et al., 2009; Ghasemzadeh et al., 2010). This activity is believed to be mainly due to their redox properties, that play a significant role in adsorbing and neutralizing free radicals by donating a hydrogen atom or an electron, chelating metal ions in aqueous solutions (Petti and Scully, 2009), quenching singlet and triplet oxygen, or decomposing peroxides (Zheng and Wang, 2001). Besides, the phenolic compounds possess multiple biological properties such as antitumor, antimutagenic, and antibacterial properties, and these activities might be related to their antioxidant activity (Shui and Leong, 2002).

Total flavonoid content

The total flavonoid content of six aromatic and medicinal plant leaves was shown in Table 3. The total phenolic content (mg/g) in methanolic extracts, expressed in quercetin equivalent (QE), varied between 1.83 ± 0.02 and 2.72 ± 0.01 . Amjad and Shafighi (2013) reported that annual and geographical climate differences, soil conditions, and pesticide or herbicide use may contribute to variations in antioxidant activity and flavonoid content of plants. The concentration of flavonoids in plant extracts depends on the polarity of solvents used in the extract

preparation (Min and Chun-Zhao, 2005). Flavonoids are a class of secondary plant metabolites with significant antioxidant and chelating properties. Besides phenolic compounds; the presence of flavonoids might also influence the antioxidant capacity. Antioxidant activity of flavonoids depends on the structure and substitution pattern of hydroxyl groups (Sharififar et al., 2009; Stanković, 2011). Flavonoids are polyphenolic compounds with low molecular mass, found in leguminous, fruits, flowers, and leaves (Harbone and Williams, 2000), having several biological activities. The polar extracts of the leaves of *Quercus ilex* were abundant in flavonoids, in particular quercetin, isorhamnetin-3-O-glucopyranoside, flavonolacylated glucosides, and the phenolic ellagic acid (Karioti et al., 2009). Quantitative analysis of the methanolic extract of the plant revealed that it is a rich source of acylated flavonoid glucosides (Karioti et al., 2010). Flavonoids as one of the most diverse and widespread groups of natural compounds are probably the most important natural phenolics. These compounds possess a broad spectrum of chemical and biological activities including radical scavenging properties (Agbor et al., 2005).

Effect of essential oils on the mortality rate of caterpillar (*Thaumetopoea pityocampa* Schiff) according to species

As shown in Table 4 that there is a highly significant effect of essential oil depending on the species of which maritime pine has the highest effect which reaches 80.55%. Then the HE of lavender and the eucalyptus have the same effect on the mortality rate of caterpillar that recorded (74.81 and 72.81%, respectively) and for juniper has a significant effect that presents 58.26% while cypress the weakest effect for a value of 40.49%. According to previous results, the insecticidal effect by fumigation of essential oils depends on the plant material used within the same plant species or even from one species to another, the source of the latter as well as the species of insect and its evolutionary stage. This was also demonstrated by Tunç et al. (2000); Chiasson et al.

Table 4. Effect of essential oils on the mortality rate of the caterpillar (*Thaumetopoea pityocampa*)

Species	Larvae mortality rate in %
<i>Pinus pinaster</i>	80.55 ^a
<i>Lavandulastoechas</i>	74.81 ^{ba}
<i>Eucalyptus camaldulensis</i>	72.77 ^{ba}
<i>Pinus pinea</i>	66.66 ^{bc}
<i>Juniperus oxycedrus</i>	58.26 ^c
<i>Cupressus sempervirens</i>	40.49 ^d

(2001); Choi et al. (2003); Sedy and Koschier (2003); Negahban et al. (2007).

CONCLUSION

The pine processionary caterpillar is a real danger to the forest ecosystem and more specifically to pine forests. This insect continues to damage most of the Northwestern pine species. According to the pine stand infestations map in the regions studied, it is depicted that maritime pine and gable pine are the most affected species by this primary insect: they recorded serious defoliation in the stand and they marked noteworthy growth losses. Analysis of the physicochemical properties of the six species in the laboratory revealed that the difference in physico-chemical composition, secondary metabolites, composition affects the insect's preference. However, the lipid composition seems to have no effect on the pine processionary caterpillar. Study of the interaction of adult larvae and essential oils revealed that maritime pine has an important effect in controlling the caterpillar. The pine processionary caterpillar cycle lasts for a year: it begins in September when the larvae are at instar 1 and ends in August. This phase is spread out between February and March.

Phytochemical analysis of the essential oil of some species has shown encouraging results in order to prepare an effective biological product to control the pine processionary caterpillar and to minimize the pine damage by this insect.

REFERENCES

- Agbor** GA, Oben JE, Ngogang JY, Xinxing C, Vinson JA, **2005**. Antioxidant capacity of some herbs/spices from cameroon: a comparative study of two methods. *J Agric Food Chem*, 53: 6819–6824.
- Amjad** L, **Shafighi** M, **2013**. Evaluation of Antioxidant Activity, Phenolic and Flavonoid Content in *Punicagranatum* var. Isfahan Malas Flowers. *Int J Agric Crop Sci*, 5(10): 1133-1139.
- Andarwulan** N, Batari R, Sandrosari DA, Bolling B, Wijaya H, **2010**. Flavonoid content and antioxidant activity of vegetable from Indonesian. *Food Chem*, 121: 1231-1235.
- Ayadi** S, Jerribi C, Abderrabba M, **2011**. Extraction and a study of essential oils of *Rosmarinus officinalis* collected from three different regions in Tunisia. *J Soc Alger Chim*, 21(1): 25-33.
- Bailen** M, Julio LF, Diaz CE, Sanz J, Martínez-Díaz RA, Cabrera R, Burillo J, Gonzalez-Coloma A, **2013**. Chemical composition and biological effects of essential oils from *Artemisia absinthium* L. cultivated under different environmental conditions. *Ind Crop Prod*, 49: 102–107.
- Bakkali** F, Averbeck S, Averbeck D, Idaomar M, **2008**. Biological effects of essential oils – a review. *Food Chem Toxicol*, 46: 446–475.
- Benhammou** N, Atik Bekkara F, Kadifkova Panovsk T, **2009**. Antioxidant activity of methanolic extracts and some bioactive compounds of *Atriplex halimus*. *C. R. Chimie*, 12: 1259–1266.
- Benincá** JP, Dalmarco JB, Pizzolatti MG, Frode TS, **2011**. Analysis of the anti-inflammatory properties of *Rosmarinus officinalis* L. in mice. *Food Chem*, 124: 468–475.
- Bligh** EG, **Dyer** WJ, **1959**. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol*, 37: 911–917.
- Braca** A, De Tommasi N, Di Bari L, Pizza C, Politi M, Morelli I, **2001**. Antioxidant Principles from *Bauhinia tarapotensis*. *J Nat Prod*, 64: 892–895.
- Chiasson** H, Be' Langer A, Bostanian N, Vincent C, Poliquin A, **2001**. Acaricidal Properties of *Artemisia absinthium* and *Tanacetum vulgare* (Asteraceae) Essential Oils Obtained by Three Methods of Extraction. *J Econ Entomol*, Vol. 94, no. 1.
- Choi** JH, Murray AS, Cheong CS, Hong DG, Chang HW, **2003**. The resolution of stratigraphic inconsistency in the luminescence ages of marine terrace sediments from Korea. *Quat Sci Rev*, 22: 1201–1206.
- Derwich** E, Manar A, Benziane Z, Boukir A, **2010**. GC/MS Analysis and In vitro Antibacterial Activity of the Essential Oil Isolated from Leaf of *Pistacialentiscus* Growing in Morocco. *World Appl Sci J*. 8: 1267-1276.
- Dewanto** VX, Wu K, Adom K, Liu RH, **2002**. Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *J Agric Food Chem*, 50: 3010–3014.
- Djeridane** A, Yous M, Nadjemi B, Boutassouna D, Stocker P, Vidal N, **2006**. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem*, 97: 654-660.
- Ebrahimi** NS, Hadian J, Mirjalili MH, Sonboli A, Yousefzadi M, **2008**. Essential oil composition and antimicrobial activity of *Thymus caramanicus* at different phenological stages. *Food Chem*, 110: 927-931.
- Eghdami** A, **Sadeghi** F, **2010**. Determination of Total Phenolic and Flavonoids Contents in Methanolic and Aqueous Extract of *Achillea Millefolium*. *Org Chem J*, 2: 81-84.
- FAO**, **2010**. Global network on integrated soil management for sustainable use of salt-affected soils. - Rome, Italy: FAO Land and Plant Nutrition Management Service.
- Gandour** M, Tamaalli W, Trabelsi N, Hessini K, Sebei K, Debez A, Abdely C, **2011**. How to optimize the seed and seed-oil production in the cash crop halophyte *Cakilemaritima*? *J Med Plants Res*, 5: 5982-5987.
- Ghasemzadeh** A, Jaafar Hawa ZE, Rahmat A, **2010**. Antioxidant Activities, Total Phenolics and Flavonoids Content in Two Varieties of Malaysia Young Ginger (*Zingiber officinale* Roscoe). *Molecules*, 15: 4324-4333.
- Habibi** H, Mazaheri D, Majnoon-Hassani N, **2006**. Investigating the effects of altitudes on the essential oil and herbal compositions of *Thymus kotschyanus* in Taleghan. *J Res Construct*, 19(4): 2-10.

- Harbone JB, Williams CA, 2000.** Advances in flavonoid research since 1992. *Phytochemistry*, 55: 481–504.
- Hashempour A, Ghazvini RF, Bakhshi D, Sanam SA, 2010.** Fatty acids composition and pigments changing of virgin olive oil (*Olea europea* L.) in five cultivars grown in Iran. *Aust J Crop Sci*, 4(4): 258-263.
- Hosni K, Jemli M, Dziri S, M'rabet Y, Ennigrou A, Casabianca H, Vulliet E, Ben Brahim N, Sebei H, 2011.** Changes in phytochemical, antimicrobial and free radical scavenging of the Peruvian peppertree (*Schinus molle* L.) as influenced by fruit maturation activities. *Ind Crop Prod*, 34: 1622–1628.
- Islam MS, Yoshimoto M, Ishigure K, Okuno S, Yamakawa O, 2003.** Effect of artificial shading and temperature on radical scavenging activity and polyphenolic composition in sweet potato (*Ipomoea batatas* L.) leaves. *J Amer Soc Hort Sci*, 128: 182-187.
- Jemâa JMB, 2014.** Essential oil as a source of bioactive constituents for the control of insect pests of economic importance in Tunisia. *Med Aromat Plants*, 3(2): 158.
- Karioti A, Bilia AR, Gabbiani C, Messori L, Skaltsa H, 2009.** Proanthocyanidin glycosides from the leaves of *Quercus ilex* L. (Fagaceae). *Tetrahedron Lett*, 50: 1771–1776.
- Karioti A, Bilia AR, Skaltsa H, 2010.** *Quercus ilex* L.: A rich source of polyacylated flavonoid glucosides. *Food Chem*, 123:131–142.
- Kouyokhi TE, Naghavi MR, Alayh M, 2008.** Study of the essential oil variation of *Ferula gummosa* samples from Iran. *Chem Nat Compd*, 44: 124–126.
- Kumaran A, Karunakaran RJ, 2006.** Antioxidant and free radical scavenging activity of an aqueous extract of *Coleus aromaticus*. *Food Chem*, 97: 109–114.
- Lee KW, Kim YJ, Lee HJ, Lee CY, 2003.** Cocoa has more phenolic phytochemicals and a higher antioxidant capacity than teas and red wine. *Food Chem*, 5: 7292-7295.
- Mahdavi M, Jouri MH, Mahmoudi J, Rezazadeh F, Mahzooni-Kachapi SS, 2013.** Investigating the altitude effect on the quantity and quality of the essential oil in *Tanacetum polycephalum* Sch. Bip. *polycephalum* in the Baladeh region of Nour, Iran. *Chin J Nat Med*, 11(5): 0553–0559.
- Miliauskas G, Venskutonis PR, Van Beek TA, 2004.** Screening of radical scavenging activity of some medicinal and aromatic plant extract. *Food Chem*, 85: 231-237.
- Min G, Chun-Zhao L, 2005.** Comparison of techniques for the extraction of flavonoids from cultured cells of *Saussurea medusa* Maxim. *World J Microb Biot*, 21: 1461-1463.
- Mothana RA, Alsaïd MS, Hasoon SS, Al-Mosaiyb NM, Al-Rehaily AJ, Al-Yahya MA, 2012.** Antimicrobial and antioxidant activities and gas chromatography mass spectrometry (GC/MS) analysis of the essential oils of *Ajuga bracteosa* Wall. ex Benth. and *Lavandula dentata* L. growing wild in Yemen. *J Med Plants Res*, 6: 3066–3071.
- Msaada K, Hosni K, Ben Taarit M, Hammami M, Marzouk B, 2009.** Effects of growing region and maturity stages on oil yield and fatty acid composition of coriander (*Coriandrum sativum* L.) fruit. *Sci Hortic*, 120: 525–531.
- Okoh OO, Sadimenko AP, Afolayan AJ, 2010.** Comparative evaluation of the antibacterial activities of the Essential Oils of *Rosmarinus officinalis* L. obtained by hydrodistillation and solvent free microwave extraction methods. *Food Chem*, 120: 308 - 312.
- Oliveira GL, Moreira DL, Mendes ADR, Guimarães EF, Figueiredo LS, Kaplan MAC, Martins ER, 2013.** Growth study and essential oil analysis of *Piper aduncum* from two sites of Cerrado biome of Minas Gerais State, Brazil. *Rev Bras Farmacogn*, 23: 743-753.
- Petti S, Scully C, 2009.** Polyphenols, oral health and disease: a review. *J Dent*, 37: 413-423.
- Ravi R, Prakash M, Bhat KK, 2007.** Aroma Characterization of coriander (*Coriandrum sativum* L.) oil samples. *Eur Food Res Technol*, 225: 367-374.
- Sangun MK, Aydin E, Timur M, Karadeniz H, Caliskan M, Ozkan A, 2007.** Comparison of chemical composition of the essential oil of *Laurus nobilis* L. leaves and fruits from different regions of Hatay, Turkey. *J Environ Biol*, 28(4): 731-733.
- Shariffif F, Dehghn-Nudeh G, Mirtajalini M, 2009.** Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chem*, 112: 885-888.
- Shui G, Leong LP, 2002.** Separation and determination of organic acids and phenolic compounds in fruit juices and drinks by high-performance liquid chromatography. *J Chromatogr A*, 977: 89 - 96.
- Singleton VL, Orthofer R, Lamuela-Raventos RM, 1999.** Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin Ciocalteu reagent. *Meth Enzymol*, 299: 152-178.
- Skoula M, Abbas JE, Johnson CB, 2000.** Genetic variation of volatiles and rosmarinic acid in populations of *Salvia fruticosa* mill growing in Crete. *Biochem Syst Ecol*, 28: 551- 561.
- Stanković M, S, 2011.** Total phenolic content, flavonoid concentration and antioxidant activity of *Marrubium peregrinum* L. extracts. *Kragujevac J Sci*, 33: 63-72.
- Tabassum N, Vidyasagar GM, 2013.** Antifungal investigations on plant essential oils. A review. *Int J Pharm Pharm Sci*, 5(2): 19-28.
- Tosun M, Ercisli S, Sengul M, Ozer H, Polat T, 2009.** Antioxidant properties and total phenolic content of eight *Salvia* species from Turkey. *Biol Res*, 41: 175-181.
- Tunç İ, Berger BM, Erler F, Dağlı F, 2000.** Ovicidal activity of essential oils from five plants against two stored-product insects. *J Stored Prod Res*, 36: 161-168.
- Uzun B, Ulger S, Cagiran IM, 2002.** Comparison of determinate and indeterminate types of sesame for oil content and fatty acid composition. - *Turkish Journal of Agriculture and Forestry* 26: 269-274.
- Wei A, Shibamoto T, 2010.** Antioxidant Lipoxygenase Inhibitory Activities and Chemical Compositions of Selected Essential Oil. *J Agric Food Chem*, 58: 7218-7225.
- Yazdani D, Jamshidi A, Mojab F, 2002.** Comparing the essential amount and the menthole in peppermint planted in different regions of the country. *Quart J Iran Aromat Herbs*, 1(3): 73-78.
- Zaouali Y, Chograni H, Trimech R, Boussaid M, 2013.** Changes in essential oil composition and phenolic fraction in *Rosmarinus officinalis* L. var. *typicus* Batt. organs during growth and incidence on the antioxidant activity. *Ind Crops Prod*, 43: 412–419.
- Zheng W, Wang YS, 2001.** Antioxidant Activity and Phenolic Compounds in Selected Herbs. - *J. Agric. Food Chem.* 49: 5165-5170.

Citation: Hasnaoui F, Zouaoui I, Dallali S, Dibouba F, 2020. Insecticidal effects of essential oils from six aromatic and medicinal plants on the pine processionary caterpillar (*Thaumetopoea pityocampa* Schiff). *Adv Med Plant Res*, 8(4): 81-88.
